



Synthesis & characterization of N-thiadiazolyl thiazolidinone derivatives

Harika MS¹, Sudha BN²

¹Creative Educational Society
College of Pharmacy, Kurnool,
Andhra Pradesh, India, ²Department
of Chemistry, JNTU-Ananthapur
University, Andhra Pradesh, India

Address for Correspondence
M.Sai Harika,
E-mail : harika.rubi@gmail.com

Received : 14-04-2014
Review completed : 21-05-2014
Accepted : 31-05-2014

ABSTRACT

Thiazolidine-4-ones containing thiazole moiety. It had been synthesized by 6-aminocoumarin, isatin, primary amines, and aromatic aldehydes. Present article is sincere attempt to synthesis of a series of {2-phenyl-3-(5-phenyl-1,3,4-thiadiazol-2-yl)-1,3-thiazolidin-4-one derivatives} (3a-3e). The synthetic approach involves the cyclocondensation of the appropriate Schiff bases with α -mercaptoalkanoic acid in glacial acetic acid. Here suitably substituted Schiff's bases (2a-2e) obtained by the reaction of appropriate aminothiadiazole(1) derivative with different substituted aromatic aldehydes. The synthesized complexes have been identified by their melting point, R_f values and interpreted by their spectral studies like IR, ¹H NMR & Mass.

Key words: Thiosemicarbazide, Thiazolidinone, Glacial acetic acid, Aromatic

Access this article online

QR Code



Website:
www.ijrpsonline.com

INTRODUCTION

Heterocyclic compounds occupy a central position among those molecules that makes life possible. The chemistry of heterocyclic compounds has been an interesting field of study for a long time. 1,3,4-thiadiazole¹ derivatives possess interesting biological activities²⁻¹⁰ probably conferred to them due to strong aromaticity of the ring system which leads to great in vivo stability and generally, a lack of toxicity for higher vertebrates, including humans when diverse functional group that interact with biological receptor are attached to aromatic ring¹¹. Compounds carrying the thiazolidinone ring have been reported to demonstrate a wide range of pharmacological activities¹² which include antimicrobial activity^{13,14}, antifungal activity¹⁵, CFTR inhibitor¹⁶, antitumor^{17,18}, anti-inflammatory¹⁹, anticonvulsant²⁰. Thiazolidinone ring can be prepared by nucleophilic addition of thioglycolic/thiolactic acid to C=N double bond²¹⁻²³.

MATERIALS AND METHOD

2-Phenyl-3-(5-Phenyl-1,3,4-Thiadiazol-2-yl)-1,3 Thiazolidin-4-one derivatives are synthesized using three steps:

synthesis of phenyl thiadiazolamine, synthesis of schiff's base, and synthesis of thiazolidinone derivatives. Melting points were determined by open capillaries on tempo apparatus. The completion of reaction and purity was checked by HPTLC pre-coated plates. IR spectra was recorded on spectrometer using KBr pressed pellet technique. ¹H NMR spectra was recorded on BRUKER 300 (MHZ) NMR spectrometer.

SYNTHESIS OF 5-PHENYL-1,3,4-THIAZOL-2-AMINE (1)

Thiosemicarbazide (0.025 mol) was suspended in a 1,4-dioxane (30ml) and stirred with the addition of benzoic acid (0.03mol). Then POCl₃ was added at 0°C. The mixture was heated at 85°C for 5hrs and left to room temperature. Evaporate the solvent and pour that solid mass into ice cold water (30ml) with good stirring. Basify the mixture to pH-9 by the addition of 40% NaOH solution. Cool room temperature, the precipitate was filtered and give washings with cold water to remove all the coloured matter, the product was dried and collect it & recrystallized from ethanol. Melting point of the compound is 211-213°C with 89.6% yield.

SYNTHESIS OF 5-PHENYL-N-[PHENYLMETHYLIDENE]-1,3,4-THIADIAZOL-2-AMINE (2a):

To 1 equivalent of compound **1** add 5 equivalent of benzaldehyde in ethanol(25ml) with 0.5ml acetic anhydride and refluxed for 8hrs and cooled. The reaction was cooled and poured in a thin stream with stirring into crushed ice contained in a 500ml beaker. The product was filtered and dried. The product was purified by recrystallization from chloroform. Yield 34.05%, melting point 207-211°C

SYNTHESIS OF N-[[4-(DIMETHYLAMINO)PHENYL]METHYLIDENE]-5-PHENYL-1,3,4-THIADIAZOL-2-AMINE (2b):

To 1 equivalent of compound **1** add 5 equivalent of 4-(dimethylamino) benzaldehyde in ethanol (25ml) with 0.5ml acetic anhydride and refluxed for 8hrs and cooled. The solid thus obtained was collected through filtration if not, pour the reaction mixture into crushed ice and collect the product. Recrystallization from chloroform. Yield 54.7%, melting point 253-257°C.

SYNTHESIS OF 4-[[5-PHENYL-1,3,4-THIADIAZOL-2-YL]IMINO]METHYL}PHENOL (2c):

To 1 equivalent of compound **1** add 5 equivalent of 4-hydroxy benzaldehyde in ethanol(25ml) with 0.5ml acetic anhydride and refluxed for 8hrs and cooled. The solid thus obtained was collected through filtration if not, pour the reaction mixture into crushed ice and collect the product. Recrystallization from chloroform. Yield 39.8%, melting point 243-251°C.

SYNTHESIS OF N-[(4-NITROPHENYL)METHYLIDENE]-5-PHENYL-1,3,4-THIADIAZOL-2-AMINE (2d):

To 1 equivalent of compound **1** add 5 equivalent of 4-nitro benzaldehyde in ethanol(25ml) with 0.5ml acetic anhydride and refluxed for 8hrs and cooled. The solid thus obtained was collected through filtration if not, pour the reaction mixture into crushed ice and collect the product. Recrystallization from chloroform. Yield 42.4%, melting point 217-222°C.

SYNTHESIS OF N-[(4-METHOXYPHENYL)METHYLIDENE]-5-PHENYL-1,3,4-THIADIAZOL-2-AMINE (2e):

To 1 equivalent of compound **1** add 5 equivalent of 4-methoxy benzaldehyde in ethanol (25ml) with 0.5ml acetic anhydride and refluxed for 8hrs and cooled. The solid thus obtained was collected through filtration if not, pour the reaction mixture into crushed ice and collect the product. Recrystallization from chloroform. Yield 36.2%, melting point 204-206°C.

SYNTHESIS OF 2-(4-PHENYL-3-(5-PHENYL-1,3,4-THIADIAZOL-2YL)-1,3-THIAZOLIDIN-4-ONE (3a):

To the Schiff's base (2a) add thioglycolic acid in equimolar ratio and reflux the above mixture using glacial acetic acid as a solvent for 10-12 hrs, cool then mix the reaction mixture with sodium carbonate (10%). The

resultant neutral solid is poured into crushed ice, collect the solid after more washings with cold water. Recrystallized from ethanol. Yield 73.1%, melting point 204-209°C.

IR (cm⁻¹ KBr): C-S-C-str (688.85), C=N-str (1690), C=O-str (1735), N-CH₂-str (2864), Ar-CH Str (3037.40), Ar-C=C-Str (1558.20); **¹HNMR ppm (DMSO):** δ 7.2-7.9(H,m,ArH), δ 2.5(2H,s,CH₂), δ 6.5(H,s,CH).

SYNTHESIS OF 2-(4-DIMETHYLAMINOPHENYL-3-(5-PHENYL-1,3,4-THIADIAZOL-2YL)-1,3-THIAZOLIDIN-4-ONE (3b):

To the Schiff's base (2b) add thioglycolic acid in equimolar ratio and reflux the above mixture using glacial acetic acid as a solvent for 10-12 hrs, cool then mix the reaction mixture with sodium carbonate (10%). The resultant neutral solid is poured into crushed ice, collect the solid after more washings with cold water. Recrystallized from ethanol. Yield 54.7%, melting point 201-204°C.

IR (cm⁻¹ KBr): C-S-C-str (688.85), C=N-str (1690), C=O-str (1735), N-CH₂-str (2864), Ar-C-H Str (3037.40), Ar-C=C Str (1558.20); **¹HNMR ppm (DMSO):** δ 7.2-7.9(9H,m,ArH), δ 2.54(2H,s,CH₂), δ 5.969(H,s,CH), δ 3.010 H of N(CH₃)₂.

SYNTHESIS OF 2-(4-HYDROXYPHENYL-3-(5-PHENYL-1,3,4-THIADIAZOL-2YL)-1,3-THIAZOLIDIN-4-ONE (3c):

To the Schiff's base (2c) add thioglycolic acid in equimolar ratio and reflux the above mixture using glacial acetic acid as a solvent for 10-12 hrs, cool then mix the reaction mixture with sodium carbonate (10%). The resultant neutral solid is poured into crushed ice, collect the solid after more washings with cold water. Recrystallized from ethanol. Yield 77.7%, melting point 217-223°C.

IR (cm⁻¹ KBr): C-S-C-str (689), C=N-str (1068), C=O-str (1697), NCH₂-str(2902.98), OH-str (3550.42), Ar C=C-str (1560); **¹HNMR ppm (DMSO):** δ 7.2-8.3(9H,m,ArH) δ 2.543(2H,s,CH₂), δ 5.9(H,s,CH), δ 13.043(s,1H,OH).

SYNTHESIS OF 2-(4-NITROPHENYL-3-(5-PHENYL-1,3,4-THIADIAZOL-2YL)-1,3-THIAZOLIDIN-4-ONE (3d):

To the Schiff's base (2d) add thioglycolic acid in equimolar ratio and reflux the above mixture using glacial acetic acid as a solvent for 10-12 hrs, cool then mix the reaction mixture with sodium carbonate (10%). The resultant neutral solid is poured into crushed ice, collect the solid after more washings with cold water. Recrystallized from ethanol. Yield 54.92%, melting point 201-205°C.

IR (cm⁻¹ KBr): Thiadiazole (1066), C-S-C-str (687.30),C=N-str (1687), C=O(1817),NCH₂-str (2900.95), Ar-C-H-Str (3030.66), Ar-C=C-Str (1550.97); **¹HNMR ppm (DMSO):** δ 7.2-7.9(9H,m,ArH), δ 2.54(2H,s,CH₂), δ 5.969(H,s,CH).

SYNTHESIS OF 2-(4-METHOXYPHENYL-3-(5-PHENYL-1,3,4-THIADIAZOL-2YL)-1,3-THIAZOLIDIN-4-ONE (3e):

To the Schiff's base (2e) add thioglycolic acid in equimolar ratio and reflux the above mixture using glacial

acetic acid as a solvent for 10-12 hrs, cool then mix the reaction mixture with sodium carbonate (10%). The resultant neutral solid is poured into crushed ice, collect the solid after more washings with cold water. Recrystallized from ethanol. Yield 72.19%, melting point 227-230°C.


IR (cm⁻¹ KBr): C-S-C-str (689.59), C=N-str (1691), C=O-str (1816), N-CH₂-str (2903.40), Ar-C-H Str (3035.33), Ar-C=C Str (1558.83)

Table 1. represents the substitutions of synthesized derivatives with their respective R_f values.

Compound	R	R _f value
3a	H	0.29
3b	N(CH ₃) ₂	0.38
3c	4-OH	0.29
3d	4-NO ₂	0.39
3e	4-OCH ₃	0.31

Solvent system to monitor TLC(precoated)= *n*-Hexane:Ethyl acetate(8:2)

Scheme



RESULTS AND DISCUSSION

In our present investigation, different derivatives of 2-phenyl-3-(5-phenyl-1,3,4-thiadiazol-2-yl)-1,3-thiazolidin-4-one (3a-3j). In synthetic scheme benzoic acid on reaction with thiosemicarbazide yielded 5-phenyl-1,3,4-thiadiazol-

2-amine **1** which on reaction with different aromatic aldehydes afforded 5-phenyl-N-[(E)-phenylmethylidene]-1,3,4-thiadiazol-2-amine derivatives (**2a-2e**). The compounds **2(a-e)** on treatment with thioglycolic acid in presence of ZnCl₂ gave 2-phenyl-3-(5-phenyl-1,3,4-thiadiazol-2-yl)-1,3-thiazolidin-4-one (**3a-3e**). The structures of the compounds were confirmed by melting points, UV-Visible spectroscopy, IR spectra in ¹H NMR and MASS spectral data. The structures of compounds (**3a-3e**) were confirmed on basis of spectral data. IR spectrum showed absorption peaks at 688 cm⁻¹ for the C-S-C of thiadiazole and absorption peaks around 1600cm⁻¹ and 2900 cm⁻¹ for the C=O & NCH₂S of thiazolidinone respectively. The ¹HNMR spectra exhibiting multiple peaks attributed to the proton at δ 7.2- 8 indicating the presence of aromatic proton while siglets at δ 2.5(CH₂), δ 5.9 (CH) indicating the thiazolidinone linkage.

REFERENCES

- Mishra G, Singh AK, Jyoti K. Review article on 1, 3, 4-Thiadiazole derivatives and it's Pharmacological activities. International Journal of ChemTech Research. 2011; 3: 1380-93.
- Swamy SN, Priya BS, Prabhswamy B, Doreswamy BH, Prasad JS, Rangappa KS. Synthesis of pharmaceutically important condensed heterocyclic 4,6-disubstituted-1,2,4-triazolo-1,3,4-thiadiazole derivatives as antimicrobials; European Journal of Medicinal Chemistry. 2006; 41: 531-38.
- Aly A and El-sayed R. Synthesis and Biological Activity of New 1,3,4-Thiadiazole Derivatives. Chem. Pap. 2006; 60(1): 50-60.
- Hadzadeh F, Vosooghi R. Synthesis of 5-(5-Amino-1,3,4-thiadiazol-2-yl)-2-imidazolylthio]acetic acids. Journal of Heterocyclic Chemistry. 2008; 45: 1-3.
- Salimon J, Salih N, Hameed A, Ibraheem H, & Yousif E. Synthesis and Antibacterial Activity of Some New 1, 3, 4-Oxadiazole and 1, 3, 4-Thiadiazole Derivatives. Journal of applied sciences research. 2010; 6(7): 866-70.
- Dogan HN, Duran A, Rollas S, Sener G, Uysal M K, & Gulen D. Synthesis of new 2, 5-disubstituted-1, 3, 4-thiadiazoles and preliminary evaluation of anticonvulsant and antimicrobial activities. Bioorganic & medicinal chemistry 2002; 10(9): 2893-98
- Cai-Jun C, Song BA, Yang S, Guang-Fang Xu et al. Synthesis and antifungal activities of 5-(3,4,5-trimethoxyphenyl)-2-sulfonyl-1,3,4- thiadiazole and 5-(3,4,5-trimethoxyphenyl)-2-sulfonyl-1,3,4-oxadiazole derivatives. Bioorganic & Medicinal Chemistry 2007; 15: 3981-89.
- Gupta A, Mishra P, Kashaw SK, Kashaw V, Stables JP. Synthesis of 3-aryl amino/amino-4-aryl-5-imino-D2-1,2,4-thiadiazoline and evaluated for anticonvulsant activity. European Journal of Medicinal Chemistry. 2008; 43: 749-54.
- Shashikant RP, Prajact K, Dighe NS, Nirmal AS, Deepak S M, Smita KP, Aarti VD. Synthesis and biological evaluation of some 1, 3, 4-thiadiazoles.

- Journal of Chemical and Pharmaceutical Research. 2009; 1(1):191-98.
10. Rajesh Sharma, Jitendra S, Chatuvedi SC. 2-Amino-5-sulfanyl- 1,3,4-thiadiazoles: A new series of selective cyclooxygenase-2 inhibitors. *Acta Pharm.* 2008; 58: 317–26.
 11. Sevgi K, Ufuk C, Bilgehan BD, Ezequiel M et al. Synthesis and cytotoxic activity of some 1,2,4-triazoline-3-thione and 2,5-disubstituted-1,3,4-thiadiazole derivatives. *Marmara Pharmaceutical Journal.* 2010; 14: 84-90.
 12. Verma, Amit, Saraf KS. 4-Thiazolidinone—a biologically active scaffold. *European journal of medicinal chemistry.* 2008; 43(5): 897-905.
 13. Bozdag-Dundar, Oya, et al. Synthesis and antimicrobial activity of some new thiazolyl thiazolidine-2, 4-dione derivatives. *Bioorganic & medicinal chemistry.* 2007; 15(18): 6012-17.
 14. Altintas H et al., Synthesis of Mannich bases of some 2,5-disubstituted 4-thiazolidinones and evaluation of their antimicrobial activities. *Turk J Chem* 2005; 29:425- 35.
 15. Rao et al., Synthesis of new 2,3-diaryl-1,3-thiazolidin-4-ones as anti-HIV agents. *IL Farmaco* 2004; 59: 33-39.
 16. Alessandro Taddei et al., Altered channel gating mechanism for CFTR inhibition by a high affinity thiazolidinone blocker. *FEBS Letters.* 2004; 558: 52-56.
 17. Abhinit, Mulay, Mangesh G, Pratima NA. Exploring potential of 4-thiazolidinone: a brief review. *International Journal of Pharmacy and Pharmaceutical Sciences.* 2009; 1(1): 47-64.
 18. Shay JW and Bacchetti S. A survey of telomerase activity in human cancer. *European journal of cancer.* 1997; 33(5): 787-91.
 19. Geronikaki A, Lagunin A, Poroikov V et al. Computeraided discovery of antiinflammatory thiazolidinones with dual cyclooxygenase/lipoxygenase inhibition. *Journal of Medicinal Chemistry.* 2008; 51(6): 1601-09.
 20. Ulusoy N, Ergenc N, Ekinc AC, & Ozer H. Synthesis and anticonvulsant activity of some new arylidenehydrazides and 4-thiazolidinones. *Monatshefte fur Chemie/Chemical Monthly.* 1996; 127(11): 1197-1202.
 21. Brown FC, Jones RS, & Kent M. The aldol condensation with 2, 3-diphenyl-4-thiazolidinone. *Canadian Journal of Chemistry.* 1963; 41(4): 817-20.
 22. Horton DA, Bourne GT, Smyth ML, The combinatorial synthesis of Bicyclic privileged structures or privileged substructures. *Chem Rev* 2003; 103: 893.
 23. Knott EB, The electrophilic reactivity of alkoxyalkylidene derivatives of heterocyclic keto methylene compounds. *J Chem Soc* 1954; 1482.