



Simultaneous estimation of Thiocolchicoside and Hydrochlorothiazide by Q-absorption ratio method

Ghanchi AA, Maheshwari DG

Dept. of Quality Assurance, L.J
Institute of Pharmacy, Ahmadabad,
Gujrat, India

Address for Correspondence
Dr. Dilip G. Maheshwari
E-mail : dgmaheshwari@gmail.com

Received: 18-02-2015
Review completed: 23-03-2015
Accepted: 25-03-2015

Access this article online

QR Code



Website:
www.ijrpsonline.com

ABSTRACT


A simple, precise and accurate UV spectrophotometric method has been developed and validated for the Simultaneous estimation of Thiocolchicoside(TCS) and Hydrochlorothiazide(HCTZ) by UV spectrophotometric method by Q-Absorption ratio method. For development of Q-Absorption ratio method, uses the ratio of absorbance at two selected wavelengths, one which is an iso-absorptive point and other being the λ -max of one of the two components. Thiocolchicoside and Hydrochlorothiazide have shown an iso-absorptive point at 282.60 nm in methanol. The second wavelength used is 271 nm, which is λ -max of Hydrochlorothiazide in methanol. Calibration curves were linear in range of 5 - 25 μ g/mL ($r^2=0.999$) and 5-25 μ g/mL ($r^2=0.999$).

Key words: Thiocolchicoside, Hydrochlorothiazide , Q-Absorption ratio method, Methanol

INTRODUCTION

Thiocolchicoside (TCS) chemically, N-[3-(β -D-glucopyranoxyloxy)-5,6,7,9-tetrahydro-1,2-imethoxy-10-(methylthio)-9-oxobenzo [a] heptalen-7yl]acetamide. It has selective affinity for γ -amino- butyric acid (GABA) receptors and acts on the muscular contracture by activating the GABA- inhibitory pathways thereby acting as a potent muscle relaxant. Thiocolchicoside is official in IP², USP⁴. Hydrochlorothiazide(HCTZ) is diuretic and antihypertensive, which inhibits the reabsorption of sodium and calcium at the beginning of distal convoluted. It is the 3,4-dihydro derivative of chlorothiazide. It is chemically 6-chloro-3,4-dihydro-2H-1, 2, 4-benzothiazine-7-sulphonamide-1,1-dioxide. Hydrochlorothiazide is official in IP², BP³, USP⁴. Thiocolchicoside is Centrally Acting Skeletal Muscle Relaxant having Analgesic And Anti-Inflammatory Effect and Hydrochlorothiazide is Thiazide class of Diuretics Used in treatment of Hypertension and Chronic Heart Failure¹. Longer use of Hydrochlorothiazide may Cause Joint Pain, Back Pain, Muscle Pain, to overcome these Hydrochlorothiazide is given with combination of Thiocolchicoside. On detailed literature survey, it was found that Thiocolchicoside can be

estimated by spectrophotometry⁵⁻¹¹, HPLC^{12,14}, HPTLC¹³ methods individually or in combination with other drugs. The aim of the present work is to develop and validate new spectrophotometric methods for the estimation of Thiocolchicoside and Hydrochlorothiazide. The proposed method was developed and validated according to guidelines.



MATERIALS AND METHODS

Instrumentation

A double beam UV visible spectrophotometer (UV – 1800, Shimadzu) connected to Computer loaded with spectra manager software UV probe was employed with spectral band width 1 nm and wavelength accuracy of 0.3 nm with a pair of 10 mm matched quartz cells. All weights were taken on electronic balance.

Chemicals and Reagents

The bulk drug Hydrochlorothiazide was obtained from (Intas Pharmaceuticals Ltd., Ahmedabad, Gujarat) and Thiocolchicoside was obtained from (West Coast, Gota, Ahmedabad, Gujarat). Methanol was used as a solvent throughout the experimentation.

Preparation of Standard Solutions and Synthetic Mixture

(A) Preparation of Standard stock solution:

A 10 mg of standard HCTZ and TCS were weighed and transferred to 100 ml separate volumetric flasks and dissolved in methanol. The flasks were shaken and volumes were made up to mark with methanol to give a solution containing 100 µg/ml each of HCTZ and TCS. This stock solution was used to prepare further dilutions of standard solutions.

(B) Preparation of Synthetic Mixture of Hydrochlorothiazide and Thiocolchicoside:

Acenac MR is a medicine that contain two active substances, Thiocolchicoside and Hydrochlorothiazide. The synthetic mixture of Hydrochlorothiazide and Thiocolchicoside was prepared in the ratio of 1:25. Accurately weighed Hydrochlorothiazide (1 mg) and (25mg) Thiocolchicoside were transferred into 100 ml volumetric flask and dissolved in Methanol. Excipients such as such as magnesium stearate, Carmellose Sodium, Polysorbate 80 were added in this mixture and sonicated for 20 minutes. This solution was filtered through Whatman filter paper. The filtrate was diluted to the mark with Methanol. The mixture contains 10 µg/ml of Hydrochlorothiazide and 250 µg/ml of Thiocolchicoside. From this solution 1 ml transferred into a 10 ml volumetric flask and the volume was adjusted up to the mark with methanol to make final concentration of Hydrochlorothiazide 1 µg/ml and Thiocolchicoside 25 µg/ml.

(C) Selection of Wavelength for Estimation of Hydrochlorothiazide and Thiocolchicoside:

The standard stock solution of Hydrochlorothiazide and Thiocolchicoside were scanned in the range of 200 nm to 400 nm against methanol as a blank. Maximum absorbance was obtained at 271 nm and 257 nm for Hydrochlorothiazide and Thiocolchicoside. Isoabsorptive point was found at 282.60 nm.

Q- Absorption ratio method

Absorbance ratio method uses the ratio of absorbance at two selected wavelengths, one which is an isoabsorptive point and other being the λ-max of one of the two components. From the overlay spectra of two drugs, it is evident that HCTZ and TCS show an isoabsorptive point at 282.60 nm. The second wavelength used is 271 nm,

which is the λ-max of HCTZ. Working standard solutions having concentration range 5-25 µg/ml for HCTZ and TCS were prepared in methanol and the absorbance at 271 nm and 282.60 nm were measured and absorptivity coefficients were calculated using calibration curve. The concentration of two drugs in the mixture can be calculated using following equations.

$$CX = [(QM-QY) / (QX-QY)] \times A1/ax1$$

$$CY = [(QM-QX) / (QY-QX)] \times A2/ay1$$

Where,

Cx and Cy= Concentration of Hydrochlorothiazide and Thiocolchicoside

Qx= The ratio of absorptivity of Thiocolchicoside at 271 and 282.60 nm

Qy= the ratio of absorptivity of Hydrochlorothiazide at 271 and 282.60 nm

QM = A2/A1, **QX** = ax2 / ax1 and **QY** = ay2 / ay1

A1 and A2= Absorbance of mixture at 271 nm and 282.60nm

ax1 and ay1 = Absorptivities of HCTZ and TCS at 271 nm

ax2 and ay2 = Absorptivities of HCTZ and TCS at 282.60 nm

Method Validation

UV spectrophotometric method developed was validated according to International Conference on Harmonization (ICH) guidelines for validation of analytical procedures. The Analytical method was validated for the parameters like linearity, accuracy, system precision, intra-day precision, inter-day precision/ intermediate precision/ ruggedness, robustness, limit of detection (LOD) and limit of quantitation (LOQ).

Calibration Curve for Hydrochlorothiazide and Thiocolchicoside

For Hydrochlorothiazide: An aliquots of stock solution of Hydrochlorothiazide (0.5, 1.0, 1.5, 2.0 and 2.5 ml) were pipette out in five different 10 ml volumetric flasks and further diluted to attain concentration of about 5, 10, 15, 20 and 25 µg/ml respectively. Graph of Absorbance Vs. Concentration was plotted.

For Thiocolchicoside: An aliquots of stock solution of Thiocolchicoside (0.5, 1.0, 1.5, 2.0 and 2.5 ml) were pipette out in five different 10 ml volumetric flasks and further diluted to attain concentration of about 5, 10, 15, 20 and 25 µg/ml respectively. Graph of Absorbance Vs. Concentration was plotted.




Figure.1: Overlain UV Spectrum of Thiocolchicoside and Hydrochlorothiazide (each 15µg/mL)




Figure. 2: “Overlain spectra of Hydrochlorothiazide (5-25 µg/ml)”




Figure.3:“Overlain spectra of Thicolchicoside (5 - 25 µg/ml)”

Table 1: “Calibration data at Isoabsorptive Point (282.60 nm)”

Hydrochlorothiazide (282.60 nm)			Thicolchicoside (282.60 nm)		
Conc. (µg/ml)	Mean Absorbance ± SD	% RSD	Conc. (µg/ml)	Mean Absorbance ± SD	% RSD
5	0.105 ± 0.0010	0.9523	5	0.105 ± 0.0010	0.9433
10	0.207 ± 0.0015	0.7367	10	0.207 ± 0.0015	0.7211
15	0.315 ± 0.0020	0.6594	15	0.314 ± 0.0025	0.7911
20	0.422 ± 0.0020	0.4925	20	0.421 ± 0.0026	0.6132
25	0.526 ± 0.0015	0.2902	25	0.528 ± 0.0035	0.6591

Table 2: “Calibration data at λmax (271 nm)”

Hydrochlorothiazide (271 nm)			Thicolchicoside (271 nm)		
Conc. (µg/ml)	Mean Absorbance ± SD	% RSD	Conc. (µg/ml)	Mean Absorbance ± SD	% RSD
5	0.352 ± 0.0020	0.5649	5	0.124 ± 0.0015	1.2096
10	0.691 ± 0.0040	0.5755	10	0.235 ± 0.0020	0.8438
15	1.028 ± 0.0055	0.5324	15	0.359 ± 0.0026	0.7202
20	1.365 ± 0.0070	0.5109	20	0.485 ± 0.0030	0.6160
25	1.719 ± 0.0085	0.4918	25	0.582 ± 0.0035	0.5982




Figure.4: “Calibration Curve of Hydrochlorothiazide and Thicolchicoside (5 – 25 µg/ml) at 271 nm”




Figure. 5: “Calibration Curve of Hydrochlorothiazide and Thicolchicoside (5 – 25 µg/ml) at 282.60 nm”

ACCURACY

To study the accuracy of the proposed methods, recovery studies were carried out by standard edition method at three different levels (80%, 100%, 120% of the test concentrations as per ICH guidelines). A known amount of drug was added and percentage recoveries were calculated. The results of recovery studies were satisfactory.

Table 3: “Accuracy data”

Drug	Amount Taken (µg/ml)	Amount Added (µg/ml)	Total amount of Drug (µg/ml)	Amount of Drug Recovered (µg ± SD)	% Recovery ± SD
Thicolchicoside	10	8	18	17.87 ± 0.0035	99.27 ± 1.017
	10	10	20	19.98 ± 0.0040	99.9 ± 0.678
	10	12	22	22.41 ± 0.0055	101.86 ± 1.340
Hydrochlorothiazide	10	8	18	17.96 ± 0.0045	99.77 ± 0.750
	10	10	20	20.13 ± 0.0052	100.65 ± 0.900
	10	12	22	21.89 ± 0.0040	99.5 ± 0.8407

PRECISION

Precision of the method was measured in terms of the interday and intraday precision and repeatability. The experiment was repeated for three times in a day using three different concentrations (3×3) for intraday and three different concentrations on three consecutive days for interday precision. The experiment was repeated for six times for repeatability.

Table 4: “Intraday Precision Data (n=3) for Hydrochlorothiazide”

Hydrochlorothiazide (282.60 nm)			Hydrochlorothiazide (271 nm)		
Conc. (µg/ml)	Mean Absorbance ± SD	%RSD	Conc. (µg/ml)	Mean Absorbance ± SD	%RSD
10	0.209 ± 0.0025	1.2022	10	0.698 ± 0.0035	0.5026
15	0.317 ± 0.0030	0.9463	15	1.034 ± 0.0058	0.5609
20	0.423 ± 0.0025	0.5940	20	1.374 ± 0.0090	0.6550

Table 5: “Interday Precision Data (n=3) for Hydrochlorothiazide”

Hydrochlorothiazide (282.60 nm)			Hydrochlorothiazide (271 nm)		
Conc. (µg/ml)	Mean Absorbance ± SD	%RSD	Conc. (µg/ml)	Mean Absorbance ± SD	%RSD
10	0.209 ± 0.0030	1.4570	10	0.696 ± 0.0043	0.6178
15	0.317 ± 0.0035	1.1055	15	1.035 ± 0.0066	0.6376
20	0.424 ± 0.0035	0.8276	20	1.375 ± 0.0100	0.7272

Table 6: “Repeatability Data (n=6) for Hydrochlorothiazide”

Hydrochlorothiazide (282.60 nm)			Hydrochlorothiazide (271 nm)		
Conc. (µg/ml)	Mean Absorbance ± SD	%RSD	Conc. (µg/ml)	Mean Absorbance ± SD	%RSD
15	0.318 ± 0.0019	0.6200	15	1.034 ± 0.0046	0.4452

Table 7: “Intraday Precision Data (n=3) for Thiocolchicoside”

Thiocolchicoside (282.60 nm)			Thiocolchicoside (271 nm)		
Conc. (µg/ml)	Mean Absorbance ± SD	%RSD	Conc. (µg/ml)	Mean Absorbance ± SD	%RSD
10	0.209 ± 0.0025	1.1961	10	0.237 ± 0.0025	1.0548
15	0.317 ± 0.0026	0.8201	15	0.363 ± 0.0045	1.2396
20	0.423 ± 0.0030	0.7092	20	0.489 ± 0.0040	0.8179

Table 8: “Interday Precision Data (n=3) for Thiocolchicoside”

Thiocolchicoside (282.60 nm)			Thiocolchicoside (271 nm)		
Conc. (µg/ml)	Mean Absorbance ± SD	%RSD	Conc. (µg/ml)	Mean Absorbance ± SD	%RSD
10	0.210 ± 0.0032	1.5238	10	0.239 ± 0.0041	1.7154
15	0.318 ± 0.0040	1.2578	15	0.364 ± 0.0055	1.5109
20	0.424 ± 0.0035	0.8254	20	0.491 ± 0.0065	1.3238

Table 9: “Repeatability Data (n=6) for Thiocolchicoside”

Thiocolchicoside (282.60 nm)			Thiocolchicoside (271 nm)		
Conc. (µg/ml)	Mean Absorbance ± SD	%RSD	Conc. (µg/ml)	Mean Absorbance ± SD	%RSD
15	0.318 ± 0.0033	1.037	15	0.365 ± 0.0052	1.4246

Limit of Detection (LOD):

Limit of detection can be calculated using following equation as per ICH guidelines.

$$LOD = 3.3 \times (\sigma / S)$$

Where, σ = Standard deviation of the Y intercept of calibration curve

S = Mean slope of the corresponding calibration curve

Limit of Quantification (LOQ):

Limit of quantification can be calculated using following equation as per ICH guidelines.

$$LOQ = 10 \times (\sigma / S)$$

Where, σ = Standard deviation of the Y intercept of calibration curve

S = Mean slope of the corresponding calibration curve

Table 10: “LOD and LOQ data”

Drug Name	LOD(µg/ml)	LOQ(µg/ml)
Thiocolchicoside	0.210	0.630
Hydrochlorothiazide	0.248	0.744

Table 11: “Analysis of synthetic mixture by Q – absorption ratio method”

Drug Name	Label Claim (mg)	Amt. in Synthetic Mix. (mg)	Total amt. obtained ± SD	% Recovery
Hydrochlorothiazide	100	25	24.87 ± 0.003	99.48 %
Thiocolchicoside	4	1	1.012 ± 0.0020	101.2 %

RESULTS AND DISCUSSION

The proposed method was based on spectrophotometric simultaneous estimation of Thiocolchicoside and Hydrochlorothiazide in UV region using Methanol as solvent. The proposed methods were found to be accurate, simple, rapid and reproducible. Thus the Proposed method can be applied in the routine analysis of the Thiocolchicoside and Hydrochlorothiazide.

Table 13: “Regression analysis data and summary of validation parameters”

Parameters	HCTZ		TCS	
	271 nm	282.60 nm	271 nm	282.60 nm
Wavelength (nm)	271 nm	282.60 nm	271 nm	282.60 nm
Beer's Law Limit (µg/ml)	5 – 25	5 – 25	5 – 25	5 – 25
Regression equation (y = mx + c)	y = 0.0682x + 0.0086	y = 0.0211x - 0.0021	y = 0.0233x + 0.0072	y = 0.0212x - 0.0003
Correlation coefficient (r ²)	0.999	0.999	0.998	0.999
Repeatability (%RSD, n = 6)	0.445	0.620	0.670	0.682
Intraday (n = 3) (%RSD)(µg/ml)	0.50 – 0.65	0.54 – 1.20	0.81 – 1.23	0.70 – 1.19
Interday (n = 3) (%RSD)(µg/ml)	0.61 – 0.72	0.82 – 1.45	1.32 – 1.71	0.82 – 1.52
LOD (µg/ml)	0.248		0.210	
LOQ (µg/ml)	0.744		0.630	
Accuracy (%)	99.27 – 101.86		99.5 – 100.65	

CONCLUSION

Simple, accurate, rapid and precise Q-Absorption ratio method was developed and validated for simultaneous estimation of both these drugs. The method utilizes easily available solvent for analysis of both drugs hence the method was also economic for estimation of HCTZ and TCS from synthetic mixture. The plot of absorbance versus respective concentration was found to be linear in the concentration range of 5-25 µg/ml for both Hydrochlorothiazide and Thiocolchicoside respectively. Accuracy of method was found between 98-102%. The precision (intra-day, inter-day and repeatability) of method was found within limits. This method can be successfully applied for the simultaneous estimation of Hydrochlorothiazide and Thiocolchicoside in synthetic mixture.

ACKNOWLEDGEMENTS

The authors are grateful to Intas Pharmaceutical Ltd. and West Coast for providing gift sample of Hydrochlorothiazide and Thiocolchicoside to carry out this work and Dr. Pundrikakshudu, Director of L.J. Institute of Pharmacy for providing all facilities for my project work.

REFERENCES

- Mohan H. Text book of Pathology. 6. Jaypee Brother's Medical Publisher: Chandigarh; 2010.
- Indian Pharmacopoeia. Government of India Ministry of Health and Family Welfare Published by Indian Pharmacopoeia Commission, Ghaziabad, 2014, Vol-II, pp 1900-1902, 2860-2862.
- British Pharmacopoeia, The Stationary Office On Behalf Of Medicines & Health Care Products Regulatory Agency, (MHRA), London, United Kingdom, 2009, 6th Edn, Vol- II, pp 374.
- United State Pharmacopoeia and National formulary. Asian Edition, The United State Pharmacopoeial Convention, Rockville, 2011, Vol-II, pp 917-919.
- Niar S, Rathode B. Simultaneous quantification of Nevibololhydrochloride and hydrochlorothiazide by first derivative UV-Spectroscopy. Der Pharmacia Lettre. 2013;5(2): 78-84.
- Patel R, Patel J. Simultaneous Equation Spectrophotometric Method for The Estimation of NebivololHCl and Hydrochlorthiazide in their Combined Dosage Form. International journal of Pharmaceutical and Applied Sciences. 2011; 2(1): 6-10.
- Harde M, Jadav S and Chaudhri P. Development and Validation of UV-Visible Spectrophotometric Methods for Simultaneous Estimation of Thiocolchicoside and Dexketoprofen in Bulk and Tablet Dosage Form. International Journal of Pharmaceutical Sciences and Drug Research. 2012; 4(2):160-163.
- Trivedi J, Chaudhari B. Spectrophotometric Methods for Simultaneous Estimation of Thiocolchicoside and Dexketoprofen Trometamol in Pharmaceutical Dosage Form. International journal for pharmaceutical Research Scholars. 2012; 1(1): 55-61.
- Patel D, Shah N. Simultaneous estimation of Aceclofenac, Thiocolchicoside and Rabepazole by uv spectrophotometer using multicomponent mode method. Inventi Rapid- Pharm Analysis & Quality Assurance. 2013;1(1): 880.
- Joshi R, Gupta K. Simultaneous UV-Spectrophotometric determination of Thiocolchicoside and Diclofenac in Pharmaceutical formulation. Der. Pharmacia. Sinica. 2010; 1(2):44-51.
- Joshi R and Gupta K. UV-Spectrophotometric Determination of Thiocolchicoside in capsule." Der. Pharma. Chemica. 2010;2(2):384-391.
- Vidyadhara S, Rao B, Koduri T. Analytical method development and validation for simultaneous estimation of Enalapril Maleate and Hydrochlorothiazide by RP-HPLC. Der. Pharm. Chemica. 2014; 6(1):217-223.
- Rajput D, Shirkhedkar A. Stability Studies of Thiocolchicoside in Bulk and Capsules using RP-HPTLC. Journal of Analytical Methods in Chemistry. 2013; 1-6.
- Pandya G and Joshi H. Development and validation of stability indicating HPLC assay method for simultaneous determination of Amlodipine Besylate, Olmesartan Medoxomil and Hydrochlorothiazide in Tablet formulation. Der. Pharmacia. Sinica. 2013; 4(2): 145-152.